Supplementary data

S1: modified CTAB DNA extraction protocol (courtesy of S. Janssens – RBG Meise)

Products

- CTAB 2x
  - 100 mM Tris-HCl
  - 1,4 M NaCl
  - 20 mM EDTA
  - 2% (w/v) CTAB
  - 1% (w/v) PVP-40
- Chloroform/isoamylalcohol 24:1
- Isopropanol (4°C)
- 70% ethanol (-20°c)
- TE buffer (pH 8,0) (4°C)
  - 10 mM Tris-HCl
  - 1 mM EDTA
- RNase A (1/10) (4°C)
  - 1 mg/mL RNase A

Protocol

1) The fly was immersed in 1 ml of 100% ethanol or CTAB solution in an Eppendorf tube
2) Pre-heat the water bath/thermoshaker 60°C
3) Shake the tube vigorously by hand for 2 min or by shaker at 5Hz for 2 min.
4) Remove the fly using forceps to a clean 1.5 ml SCREWCAP tube and add 500µl of CTAB or 100% ethanol to the new tube.
5) Shake the tube vigorously by hand for 2 min or by shaker at 5Hz for 2 min.
6) Remove the fly using forceps into a new 1.5ml clean tube and frozen at −20°C for subsequent species identification.
7) Transfer the entire volume of the first tube into the screw cap tube.
8) The tube containing the detergent and pollen was centrifuged at 13,000 rpm for 5 min
9) Remove the supernatant and dry the samples in the SPEEDVAC for 1h. (30min for ethanol).

10) Put three 3-mm stainless steel beads into each tube and disrupt the sample for 2 min at 22.5 Hz.
11) Resuspend the samples in 400 µl CTAB 2x extraction buffer and 0.5 µl (0,1%) of β-Mercaptoethanol (or 1,5µl (0,3%) might be tested) UNDER the hood
12) Vortex for 10 s
13) Put the sample in 60° C for 1h (or overnight)

14) Let the samples cool down.
15) Prepare tubes with 300 µl of Isopropanol and store in the fridge at -20°C for 30min.
16) Centrifuge the tubes at 13,000 rpm for 5 min. The ≈400 µl of supernatant was extracted to a clean 1.5 ml tube. KEEP SUPERNATANT and UNDER the hood

17) Add an equal volume (400µl) of Chloroform:Isoamylalcohol (24:1) UNDER the hood
18) Vortex vigorously for at least 5s and shake the sample for 2 min at 6Hz TWO Times.
19) Centrifuge for 15 min at 11000 rmp at 22°C.
20) Carefully transfer 400 µL of the upper aqueous phase to the isopropanol prepared tubes (0,8 v/v). CAREFUL not to disrupt the middle and bottom phase (by gently transferring the upper phase in multiple steps of 200 ul, aim for a total collected volume of ≈ 350 µl) With pollen we will only do this phase once. UNDER the hood
21) Shake by up and down (+- 50 times)

22) Store 20 min at -20°C for maximal precipitation
23) Centrifuge 10 minutes at 14000 rpm (preferably at 4°C).
24) Remove the supernatant by pouring the liquid in the isopropanol waste and place the tubes inverted for a few minutes.
25) Add 400 µL 70% cold ethanol and loosen the pellet.
26) Store 20 min at -20°C for maximal precipitation
27) Centrifuge for 10 min at 20000 rcf at 4°C.

28) Remove the supernatant by pouring the liquid in the ethanol waste and place the tubes inverted for a few minutes.
29) Put the tubes horizontal and let them air dry for around 1h. or use the speedvac for 30min.
30) Dissolve the pellet by adding 30ul TE buffer to the dried tubes.
31) Tap the pellet to dissolve it in the TE buffer
32) Add 3 µL RNase A (1/10) per tube, shake and spin down.
33) Incubate maximum 2 minutes at room temperature.
34) Store the DNA in the fridge or freezer.

S2: ANOVA on DNA concentrations and yields as obtained from pollen isolated from 50 flower flies. Test for the effects of pollen DNA extraction protocol (Qiagen vs CTAB) and preservation group (a, …, e, see Tab. 1). n.s.: not significant; ∗: P < 0.05, ∗∗: P < 0.01, *** P < 0.001. C: Cochran's test.

DNA concentrations

ANOVA

	Df	MS	F	
DNA extraction (Extr)	1	0.05642	0.6986	n.s.
Preservation (Pres-	4	1.04425	12.9309	***
Extr x Pres	4	0.53808	6.6630	***
Residual	40	0.08076		

Transformation = Fourth root

C = 0.854 n.s.

A posteriori comparisons: Extr x Pres

Qiagen extraction: EtOH RT = Et -20°C = control = CTAB-20°C = CTAB RT

CTAB extraction: control = Et -20°C = EtOH RT < CTAB-20°C < CTAB RT

DNA yield

ANOVA

	Df	MS	F	
DNA extraction (Extr)	1	0.7421	1.2837	ns
Preservation (Pres)	4	6.6469	11.4985	***
Extr x Pres	4	2.9009	5.0182	**
Residual	40	0.5781		

Transformation = Fourth root

C = 0.530 n.s.

A posteriori comparisons: Extr x Pres

Qiagen extraction: EtOH RT = Et -20°C = control = CTAB-20°C = CTAB RT

CTAB extraction: control = Et -20°C = EtOH RT < CTAB-20°C < CTAB RT

S3: Optimal annealing temperatures for each of the primer pairs tested

Marker	Forward primer	Reverse primer	Primer pair ID	Annealing Temperature	Primer sequences
matK	KIM3	KIM1	KIM	57°C	CGTACAGTACTTTTGTGTTTACGAG - ACCCAGTCCATCTGGAAATCTTGGTTC+
rbcL	rbcL-Tag-F	rbcL506	rbcL-506	58°C	ATG TCA CCA CAA ACA GAG ACT - AGGGGACGACCATACTTGTTCA
	rbcL-Tag-F	rbcL375	rbcL-375	58°C	ATG TCA CCA CAA ACA GAG ACT - GCAAATCCTCCAGACGTAGA
	rbcL-Tag-F	rbcL320	rbcL-320	61°C	ATG TCA CCA CAA ACA GAG ACT - ACCCACAATGGAAGTAAACATGT
	rbcL_230	rbcL506	rbcL-23506	56°C	CTTACCAGYCTTGATCGTTACAAAGG - AGGGGACGACCATACTTGTTCA
	rbcLF-Tag_IL	rbcLR-Tag_IL	rbcL-T	62°C	ATG TCA CCA CAA ACA GAG ACT - GA AAC GGT CTC TCC AAC GCA T
	rbcL_260	rbcL_230	rbcL-2623	58°C	GTAAAATCAAGTCCACCRCG - CTTACCAGYCTTGATCGTTACAAAGG
	rbcL-af	rbcL-ar	rbcL-A	/	ATGTCACCACAAACAGAGACTAAAGC - CTTCTGCTACAAATAAGAATCGATCTC
ITS1	ITS1-18S	ITS1-390	ITS1-390	58°C	AGTCGTAACAAGGTTTCCGT - GGGATTCTGCAATTCACACC
	ITS1-18S	ITS1-380	ITS1-380	60°C	AGTCGTAACAAGGTTTCCGT - AACTTGCGTTCAAAGACTCG
ITS2	ITS2	ITS3a	ITS2-23	53°C	ATGCGATACTTGGTGTGAAT - GACGCTTCTCCAGACTACAAT
	ITS3	ITS4	ITS2-34	56°C	GCA TCG ATG AAG AAC GCA GC - TC CTC CGC TTA TTG ATA TGC
	ITS5a	ITS4	ITS2-54	54°C	CCTTATCATTTAGAGGAAGGAG - TCCTCCGCTTATTGATATGC
	UniPlantF	UniplantR	ITS2-Uni	55°C	TGTGAATTGCARRATYCMG - CCCGHYTGAYYTGRGGTCDC

S4: Metadata sequenced samples

Specimen	Library Prep	Marker	sample group	Collection Year	taxon group	Species
MSC_HE_04	Semi-Custom	ITS2	recently collected	2022	Syrphidae	*Betasyrphus adligatus*
CMC_HC_02	Semi-Custom	ITS2	recently collected	2022	Syrphidae	*Ischiodon aegyptius*
CMC_HC_04	Semi-Custom	ITS2	recently collected	2022	Syrphidae	*Ischiodon aegyptius*
CMB_HC_08	Semi-Custom	ITS2	recently collected	2022	Syrphidae	*Ischiodon aegyptius*
CMC_HC_08	Semi-Custom	rbcL	recently collected	2022	Syrphidae	*Toxomerus floralis*
CMC_HC_02	Semi-Custom	rbcL	recently collected	2022	Syrphidae	*Ischiodon aegyptius*
CMC_HC_04	Semi-Custom	rbcL	recently collected	2022	Syrphidae	*Ischiodon aegyptius*
CMB_HC_08	Semi-Custom	rbcL	recently collected	2022	Syrphidae	*Ischiodon aegyptius*
MSC_HE_04	Semi-Custom	rbcL	recently collected	2022	Syrphidae	*Betasyrphus adligatus*
MSC_HE_06	Semi-Custom	rbcL	recently collected	2022	Syrphidae	*Betasyrphus adligatus*
MSW_HE_03	Semi-Custom	rbcL	recently collected	2022	Syrphidae	*Toxomerus floralis*
MSC_HE_06	Semi-Custom	ITS2	recently collected	2022	Syrphidae	*Betasyrphus adligatus*
MSC_HD_06	Semi-Custom	rbcL	recently collected	2022	Syrphidae	*Betasyrphus adligatus*
MSW_HD_08	Semi-Custom	rbcL	recently collected	2022	Syrphidae	*Toxomerus floralis*
MSW_HD_06	Semi-Custom	rbcL	recently collected	2022	Syrphidae	*Toxomerus floralis*
CMW_HC_A	Semi-Custom	rbcL	recently collected	2022	Syrphidae	*Toxomerus floralis*
CMW_HC_B	Semi-Custom	rbcL	recently collected	2022	Syrphidae	*Toxomerus floralis*
BM_4	Semi-Custom	ITS2	historical	1922	Hymenoptera, Apoidea	*Apis mellifera*

BM_5	Semi-Custom	ITS2	historical	1922	Hymenoptera, Apoidea	*Apis mellifera*
BM_32	Semi-Custom	ITS2	historical	1963	Hymenoptera, Apoidea	*Apis mellifera*
BM_33	Semi-Custom	ITS2	historical	1993	Hymenoptera, Apoidea	*Apis mellifera*
BM_4	Semi-Custom	rbcL	historical	1922	Hymenoptera, Apoidea	*Apis mellifera*
MSW_HE_03	Semi-Custom	ITS2	recently collected	2022	Syrphidae	*Toxomerus floralis*
BM_5	Semi-Custom	rbcL	historical	1922	Hymenoptera, Apoidea	*Apis mellifera*
BM_32	Semi-Custom	rbcL	historical	1963	Hymenoptera, Apoidea	*Apis mellifera*
BM_33	Semi-Custom	rbcL	historical	1993	Hymenoptera, Apoidea	*Apis mellifera*
VB_BB_02	Semi-Custom	ITS2	recently collected	2022	Hymenoptera, Apoidea	*Apis mellifera*
VB_BB_06	Semi-Custom	ITS2	recently collected	2022	Hymenoptera, Apoidea	*Apis mellifera*
VB_BB_08	Semi-Custom	ITS2	recently collected	2022	Hymenoptera, Apoidea	*Apis mellifera*
KBHB_BE_04	Semi-Custom	ITS2	recently collected	2022	Hymenoptera, Apoidea	*Apis mellifera*
KBHB_BE_07	Semi-Custom	ITS2	recently collected	2022	Hymenoptera, Apoidea	*Apis mellifera*
KBHB_BE_03	Semi-Custom	ITS2	recently collected	2022	Hymenoptera, Apoidea	*Apis mellifera*
VB_BB_02	Semi-Custom	rbcL	recently collected	2022	Hymenoptera, Apoidea	*Apis mellifera*
MSC_HD_06	Semi-Custom	ITS2	recently collected	2022	Syrphidae	*Betasyrphus adligatus*
VB_BB_06	Semi-Custom	rbcL	recently collected	2022	Hymenoptera, Apoidea	*Apis mellifera*
VB_BB_08	Semi-Custom	rbcL	recently collected	2022	Hymenoptera, Apoidea	*Apis mellifera*
KBHB_BE_04	Semi-Custom	rbcL	recently collected	2022	Hymenoptera, Apoidea	*Apis mellifera*
KBHB_BE_07	Semi-Custom	rbcL	recently collected	2022	Hymenoptera, Apoidea	*Apis mellifera*
KBHB_BE_03	Semi-Custom	rbcL	recently collected	2022	Hymenoptera, Apoidea	*Apis mellifera*
409 51D	Semi-Custom	ITS2	historical	1947	Hymenoptera, Apoidea	*Apis mellifera*
409 50D	Semi-Custom	ITS2	historical	1947	Hymenoptera, Apoidea	*Apis mellifera*
409 51D	Semi-Custom	rbcL	historical	1947	Hymenoptera, Apoidea	*Apis mellifera*
409 50D	Semi-Custom	rbcL	historical	1947	Hymenoptera, Apoidea	*Apis mellifera*
MSW_HD_08	Semi-Custom	ITS2	recently collected	2022	Syrphidae	*Toxomerus floralis*
MSW_HD_06	Semi-Custom	ITS2	recently collected	2022	Syrphidae	*Toxomerus floralis*
CMW_HC_A	Semi-Custom	ITS2	recently collected	2022	Syrphidae	*Toxomerus floralis*
CMW_HC_B	Semi-Custom	ITS2	recently collected	2022	Syrphidae	*Toxomerus floralis*
CMC_HC_08	Semi-Custom	ITS2	recently collected	2022	Syrphidae	*Toxomerus floralis*
MSC_HE_04	Nextera XT	ITS2	recently collected	2022	Syrphidae	*Betasyrphus adligatus*
CMC_HC_02	Nextera XT	ITS2	recently collected	2022	Syrphidae	*Ischiodon aegyptius*
CMC_HC_04	Nextera XT	ITS2	recently collected	2022	Syrphidae	*Ischiodon aegyptius*
CMB_HC_08	Nextera XT	ITS2	recently collected	2022	Syrphidae	*Ischiodon aegyptius*
CMC_HC_08	Nextera XT	rbcL	recently collected	2022	Syrphidae	*Toxomerus floralis*
CMC_HC_02	Nextera XT	rbcL	recently collected	2022	Syrphidae	*Ischiodon aegyptius*
CMC_HC_04	Nextera XT	rbcL	recently collected	2022	Syrphidae	*Ischiodon aegyptius*
CMB_HC_08	Nextera XT	rbcL	recently collected	2022	Syrphidae	*Ischiodon aegyptius*
MSC_HE_04	Nextera XT	rbcL	recently collected	2022	Syrphidae	*Betasyrphus adligatus*
MSC_HE_06	Nextera XT	rbcL	recently collected	2022	Syrphidae	*Betasyrphus adligatus*
MSW_HE_03	Nextera XT	rbcL	recently collected	2022	Syrphidae	*Toxomerus floralis*
MSC_HE_06	Nextera XT	ITS2	recently collected	2022	Syrphidae	*Betasyrphus adligatus*
MSC_HD_06	Nextera XT	rbcL	recently collected	2022	Syrphidae	*Betasyrphus adligatus*
MSW_HD_08	Nextera XT	rbcL	recently collected	2022	Syrphidae	*Toxomerus floralis*
MSW_HD_06	Nextera XT	rbcL	recently collected	2022	Syrphidae	*Toxomerus floralis*
CMW_HC_A	Nextera XT	rbcL	recently collected	2022	Syrphidae	*Toxomerus floralis*
CMW_HC_B	Nextera XT	rbcL	recently collected	2022	Syrphidae	*Toxomerus floralis*
BM_4	Nextera XT	ITS2	historical	1922	Hymenoptera, Apoidea	*Apis mellifera*
BM_5	Nextera XT	ITS2	historical	1922	Hymenoptera, Apoidea	*Apis mellifera*
BM_32	Nextera XT	ITS2	historical	1963	Hymenoptera, Apoidea	*Apis mellifera*
BM_33	Nextera XT	ITS2	historical	1993	Hymenoptera, Apoidea	*Apis mellifera*
BM_4	Nextera XT	rbcL	historical	1922	Hymenoptera, Apoidea	*Apis mellifera*
MSW_HE_03	Nextera XT	ITS2	recently collected	2022	Syrphidae	*Toxomerus floralis*
BM_5	Nextera XT	rbcL	historical	1922	Hymenoptera, Apoidea	*Apis mellifera*
BM_32	Nextera XT	rbcL	historical	1963	Hymenoptera, Apoidea	*Apis mellifera*
BM_33	Nextera XT	rbcL	historical	1993	Hymenoptera, Apoidea	*Apis mellifera*
VB_BB_02	Nextera XT	ITS2	recently collected	2022	Hymenoptera, Apoidea	*Apis mellifera*
VB_BB_06	Nextera XT	ITS2	recently collected	2022	Hymenoptera, Apoidea	*Apis mellifera*
VB_BB_08	Nextera XT	ITS2	recently collected	2022	Hymenoptera, Apoidea	*Apis mellifera*
KBHB_BE_04	Nextera XT	ITS2	recently collected	2022	Hymenoptera, Apoidea	*Apis mellifera*
KBHB_BE_07	Nextera XT	ITS2	recently collected	2022	Hymenoptera, Apoidea	*Apis mellifera*
KBHB_BE_03	Nextera XT	ITS2	recently collected	2022	Hymenoptera, Apoidea	*Apis mellifera*

VB_BB_02	Nextera XT	rbcL	recently collected	2022	Hymenoptera, Apoidea	*Apis mellifera*
MSC_HD_06	Nextera XT	ITS2	recently collected	2022	Syrphidae	*Betasyrphus adligatus*
VB_BB_06	Nextera XT	rbcL	recently collected	2022	Hymenoptera, Apoidea	*Apis mellifera*
VB_BB_08	Nextera XT	rbcL	recently collected	2022	Hymenoptera, Apoidea	*Apis mellifera*
KBHB_BE_04	Nextera XT	rbcL	recently collected	2022	Hymenoptera, Apoidea	*Apis mellifera*
KBHB_BE_07	Nextera XT	rbcL	recently collected	2022	Hymenoptera, Apoidea	*Apis mellifera*
KBHB_BE_03	Nextera XT	rbcL	recently collected	2022	Hymenoptera, Apoidea	*Apis mellifera*
409 51D	Nextera XT	rbcL	historical	1947	Hymenoptera, Apoidea	*Apis mellifera*
409 50D	Nextera XT	rbcL	historical	1947	Hymenoptera, Apoidea	*Apis mellifera*
409 51D	Nextera XT	ITS2	historical	1947	Hymenoptera, Apoidea	*Apis mellifera*
409 50D	Nextera XT	ITS2	historical	1947	Hymenoptera, Apoidea	*Apis mellifera*
MSW_HD_08	Nextera XT	ITS2	recently collected	2022	Syrphidae	*Toxomerus floralis*
MSW_HD_06	Nextera XT	ITS2	recently collected	2022	Syrphidae	*Toxomerus floralis*
CMW_HC_A	Nextera XT	ITS2	recently collected	2022	Syrphidae	*Toxomerus floralis*
CMW_HC_B	Nextera XT	ITS2	recently collected	2022	Syrphidae	*Toxomerus floralis*
CMC_HC_08	Nextera XT	ITS2	recently collected	2022	Syrphidae	*Toxomerus floralis*

S5: Detailed description of the custom pipeline for metabarcoding pollen

There are two Amplicon PCR, the first one will be done with ITS2 primers on the 24 samples of pollen DNA and a second one will be done with rbcL primers on the same 24 samples of pollen DNA.

After the first PCR Clean-up, 48 amplicon PCR samples will be processed.

1st Amplicon PCR:

Consumables

Item

Genomic DNA

Amplicon PCR Reverse Primer (20 μM)

Amplicon PCR Forward Primer (20 μM)

Taq PLATINUM

MgCl2 PLATINUM

dNTP

MQ water

96 - well 0.2 ml PCR plate 1 plate

Procedure

1 Set up the following reaction of DNA, Taq PLATINUM mix, and primers:

Volume	
5µl (or double with low quantities)	Genomic DNA (2ng/µl – 0,1ng/µl)
2,50µl	PCR Buffer(10x)
0,75µl	MgCl PLATINUM 50mM
2,50µl	dNTP 2mM
0,50µl	Primer1= LF1 20µM
0,50µl	Primer2=LR1 20µM
0,15µl	Taq PLATINUM 5 Units/µl
13,10µl	MQ H2O
Total	25 µl

2 Perform PCR in a thermal cycler using the following program:

• 94° C for 3 minutes

• 35 cycles of:

— 94° C for 30 seconds

— 55° C for 30 seconds

— 72° C for 30 seconds

• 72° C for 5 minutes

• Hold at 4° C

- Check the success of amplification with electrophoresis gel. Redo every samples than didn't work

1st PCR Clean up:

Consumables

Item Quantity Storage

MilliQ water

HighPrep™ PCR beads

Freshly Prepared 80% Ethanol (EtOH)

96 - well 0.2 ml PCR plate 1 plate

Preparation

• Bring the HighPrep™ PCR beads to room temperature.

• Prepare the plate according to plan with 20 µl and a balance

Procedure

1 Centrifuge the Amplicon PCR plate at 1,000 × g at 20° C for 1 minute to collect condensation, carefully remove seal. (Not necessary if next day)

2 Vortex the HighPrep™ PCR beads for 30 seconds to make sure that the beads are evenly dispersed. Add an appropriate volume of beads to a trough depending on the number of samples processing.

3 Using a multichannel pipette, add 20 µl of HighPrep™ PCR beads (1:1 proportion) to each well of the Amplicon PCR plate. Change tips between columns.

4 Gently pipette entire volume up and down 10 times if using a 96 - well PCR plate

5 Incubate at room temperature without shaking for 5 minutes.

6 Place the plate on a magnetic stand for 2 minutes or until the supernatant has cleared.

7 With the Amplicon PCR plate on the magnetic stand, use a multichannel pipette to remove and discard the supernatant. Change tips between samples.

8 With the Amplicon PCR plate on the magnetic stand, wash the beads with freshly prepared 80% ethanol as follows:

a Using a multichannel pipette, add 200 µl of freshly prepared 80% ethanol to each sample well.

b Incubate the plate on the magnetic stand for 30 seconds.

c Carefully remove and discard the supernatant.

10 With the Amplicon PCR plate on the magnetic stand, perform a second ethanol wash as follows:

a Using a multichannel pipette, add 200 μl of freshly prepared 80% ethanol to each sample well.

b Incubate the plate on the magnetic stand for 30 seconds.

c Carefully remove and discard the supernatant.

d Use a P20 multichannel pipette with fine pipette tips to remove excess ethanol.

11 With the Amplicon PCR plate still on the magnetic stand, allow the beads to air - dry for 10 minutes or until there is not ethanol left.

12 Remove the Amplicon PCR plate from the magnetic stand. Using a multichannel pipette, add 52.5 μl of MilliQ water to each well of the Amplicon PCR plate.

13 Gently pipette mix up and down 10 times, changing tips after each column (or seal plate and shake at 1800 rpm for 2 minutes). Make sure that beads are fully resuspended.

14 Incubate at room temperature for 2 minutes.

15 Place the plate on the magnetic stand for 2 minutes or until the supernatant has cleared.

16 Using a multichannel pipette, carefully transfer 50 μl of the supernatant from the Amplicon PCR plate to a new 96 - well PCR plate. Change tips between samples to avoid cross - contamination.

17 Using the last 2,5µl in plate, check the result of the cleaning with a gel.

SAFE STOPPING POINT

If you do not immediately proceed to *Index PCR*, seal plate

2nd Index PCR:

This step attaches dual indices and Illumina sequencing adapters using the Nextera XT Index Kit.

Consumables

Item Quantity Storage

Nextera XT Index 1 Primers (N7XX) from the Nextera XT Index kit (FC - 131 - 1001 or FC - 131 - 1002)

Nextera XT Index 2 Primers (S5XX) from the Nextera XT Index kit (FC - 131 - 1001 or FC - 131 - 1002)

Taq PLATINUM

MgCl2 PLATINUM

dNTP

MQ water

96 - well 0.2 ml PCR plate 1 plate

Microseal 'A' film

Procedure

1 Using a multichannel pipette, transfer 5 μl from each 48 well to a new 96 - well plate. The remaining 45 μl is not used in the protocol and can be stored for other uses.

2 Arrange the Index 1 and 2 primers in a rack using the following arrangements as needed:

a Arrange Index 2 primer tubes (white caps, clear solution) vertically, aligned with rows A through D. (Half a plate as the other half is going to be Nextera XT kit).

b Arrange Index 1 primer tubes (orange caps, yellow solution) horizontally, aligned with columns 1 through 12.

3 Place the 96 - well PCR plate with the 5 μl of resuspended PCR product DNA.

4 Set up the following reaction of DNA, Taq PLATINUM mix, and primers:

Volume	
5 µl	Amplicon DNA
2,50 µl	PCR Buffer(10x)
1,5 µl	MgCl PLATINUM 50mM
5,0 µl	dNTP 2mM
5 µl	Nextera XT Index Primer 1 (N7xx)
5 µl	Nextera XT Index Primer 2 (S5xx)
0,3 µl	Taq PLATINUM 5 Units/µl
34 µl	MQ H2O
Total	50 µl

5 Gently pipette up and down 10 times to mix.

6 Cover the plate with Microseal 'A'.

7 Centrifuge the plate at 1,000 × g at 20° C for 1 minute.

8 Perform PCR on a thermal cycler using the following program:

- 95° C for 3 minutes
- 8 cycles of:
  - 95° C for 30 seconds
  - 55° C for 30 seconds
  - 72° C for 60 seconds
- 72° C for 10 minutes
- Hold at 4° C

2nd PCR Clean up:

Consumables

Item	Quantity	Storage
10 mM Tris pH 8.5		
HighPrep™ PCR beads		
Freshly Prepared 80% Ethanol (EtOH)		
96 - well 0.2 ml PCR plate	1 plate	

Procedure

1 Centrifuge the Index PCR plate at 280 × g at 20° C for 1 minute to collect condensation.

2 Vortex the HighPrep™ PCR beads for 30 seconds to make sure that the beads are evenly dispersed. Add an appropriate volume of beads to a trough.

3 Using a multichannel pipette, add 56 μl of HighPrep™ PCR beads to each well of the Index PCR plate. (1:1 beads proportion for 20 µl of PCR products)

4 Gently pipette mix up and down 10 times if using a 96 - well PCR plate

5 Incubate at room temperature without shaking for 5 minutes.

6 Place the plate on a magnetic stand for 2 minutes or until the supernatant has cleared.

7 With the Index PCR plate on the magnetic stand, use a multichannel pipette to remove and discard the supernatant. Change tips between samples.

8 With the Index PCR plate on the magnetic stand, wash the beads with freshly prepared 80% ethanol as follows:

a Using a multichannel pipette, add 200 μl of freshly prepared 80% ethanol to each sample well.

b Incubate the plate on the magnetic stand for 30 seconds.

c Carefully remove and discard the supernatant.

9 With the Index PCR plate on the magnetic stand, perform a second ethanol wash as follows:

a Using a multichannel pipette, add 200 μl of freshly prepared 80% ethanol to each sample well.

b Incubate the plate on the magnetic stand for 30 seconds.

c Carefully remove and discard the supernatant.

d Use a P20 multichannel pipette with fine pipette tips to remove excess ethanol.

10 With the Index PCR plate still on the magnetic stand, allow the beads to air - dry for 10 minutes.

11 Remove the Index PCR plate from the magnetic stand. Using a multichannel pipette, add 27.5 μl of 10 mM Tris pH 8.5 to each well of the Index PCR plate.

12 If using a 96 - well PCR plate, gently pipette mix up and down 10 times until beads are fully resuspended, changing tips after each column.

13 Incubate at room temperature for 2 minutes.

14 Place the plate on the magnetic stand for 2 minutes or until the supernatant has cleared.

15 Using a multichannel pipette, carefully transfer 25 μl of the supernatant from the Index PCR plate to a new 96 - well PCR plate. Change tips between samples to avoid crosscontamination.

SAFE STOPPING POINT

If you do not immediately proceed to *Index PCR*, seal plate

Validate Library

Run 1 μl of a 1:50 dilution of the final library on a Bioanalyzer DNA 1000 chip to verify the size.

Library Quantification, Normalization, and Pooling:

Illumina recommends quantifying your libraries using a fluorometric quantification method that uses dsDNA binding dyes. Calculate DNA concentration in nM, based on the size of DNA amplicons as determined by an Agilent Technologies 2100 Bioanalyzer trace:

$$\frac{(\text{concentration in ng}/\mu\text{l})}{(660\ \text{g/mol} \times \text{average library size})} \times 10^6 = \text{concentration in nM}$$

Dilute concentrated final library using Resuspension Buffer (RSB) or 10 mM Tris pH 8.5 to 4 nM. Aliquot 5 μl of diluted DNA from each library and mix aliquots for pooling libraries with unique indices. Final volume of 480μl Depending on coverage needs, up to 96 libraries can be pooled for one MiSeq run.

S6: Detailed description of the Nextera XT DNA Library Prep Kit for metabarcoding pollen

The Nextera Kit can be used with PCR Amplicons. The PCR amplicon must be > 300 bp. Shorter amplicons can be lost during the library cleanup step.

Amplicon PCR on Genomic DNA

There is two PCR, the first one will be done with ITS2 primers on the 24 samples of pollen DNA and a second one will be done with rbcL primers on the same 24 samples of pollen DNA.

1 Set up the following reaction of DNA, Taq PLATINUM mix, and primers:

Volume	
2.5 µl	Genomic DNA (2ng/µl – 0,1ng/µl)
2,50µl	PCR Buffer(10x)
0,75µl	MgCl PLATINUM 50mM
2,50µl	dNTP 2mM
0,50µl	Primer1= LF1 20µM
0,50µl	Primer2=LR1 20µM
0,15µl	Taq PLATINUM 5 Units/µl
15,75µl	MQ H2O
Total	25 µl

2 Perform PCR in a thermal cycler using the following program:

- 95° C for 3 minutes
- 25 cycles of:
  - 95° C for 30 seconds
  - 55° C for 30 seconds
  - 72° C for 30 seconds
- 72° C for 5 minutes
- Hold at 4° C

After this first PCR, 48 samples of PCR amplicons will be processed.

Tagment Amplicon DNA

This step uses the Nextera transposome to tagment gDNA, which is a process that fragments DNA and then tags the DNA with adapter sequences in a single step.

Consumables

- ATM (Amplicon Tagment Mix)
- TD (Tagment DNA Buffer)
- NT (Neutralize Tagment Buffer)
- gDNA (0.2 ng/µl per sample)/ Amplicon DNA
- Hard-Shell 96-well PCR plate, skirted
- Microseal 'B' adhesive seals

Preparation

1. Prepare the following consumables:

Item Storage Instructions

gDNA -25°C to -15°C Thaw on ice. Invert the thawed tubes 3–5 times, and then centrifuge briefly.

ATM -25°C to -15°C Thaw on ice. Invert the thawed tubes 3–5 times, and then centrifuge briefly.

TD -25°C to -15°C Thaw on ice. Invert the thawed tubes 3–5 times, and then centrifuge briefly.

NT 15°C to 30°C Check for precipitates. If present, vortex until all particulates are resuspended.

2. Save the following tagmentation program on the thermal cycler:

- Choose the preheat lid option
- 55°C for 5 minutes
- Hold at 10°C

Procedure

1. Add the following volumes in the order listed to each well of a new Hard-Shell skirted PCR plate. Pipette to mix.
   - TD (10 µl)
   - Normalized gDNA (5 µl)
2. Add 5 µl ATM to each well. Pipette to mix.
3. Centrifuge at 280 × g at 20°C for 1 minute.
4. Place on the preprogrammed thermal cycler and run the tagmentation program. When the sample reaches 10°C, immediately proceed to step 5 because the transposome is still active.
5. Add 5 µl NT to each well. Pipette to mix.
6. Centrifuge at 280 × g at 20°C for 1 minute.
7. Incubate at room temperature for 5 minutes.

The PCR plate contains 25 µl tagmented and neutralized gDNA, all of which is used in the next step.

Amplify Libraries

This step amplifies the tagmented DNA using a limited-cycle PCR program. PCR adds the Index 1 (i7), Index 2 (i5), and full adapter sequences to the tagmented DNA from the previous step. The index adapters and Nextera PCR Master Mix are added directly to the 25 µl of tagmented gDNA from the previous step.

Consumables

- NPM (Nextera PCR Master Mix)
- Index 1 adapters (N7XX)
- Index 2 adapters (S5XX)
- TruSeq™ Index Plate Fixture
- Microseal 'A' film

Preparation

1 Prepare the following consumables:

Item Storage Instructions

Index adapters
(i5 and i7)

-25°C to -15°C Only prepare adapters being used. Thaw at room temperature for 20 minutes. Invert each tube to mix. Centrifuge briefly.

NPM -25°C to -15°C Thaw on ice for 20 minutes.

2 Save the following program on the thermal cycler:

- Choose the preheat lid option.
- 72°C for 3 minutes
- 95°C for 30 seconds
- 12 cycles of:
- 95°C for 10 seconds
- 55°C for 30 seconds
- 72°C for 30 seconds
- 72°C for 5 minutes
- Hold at 10°C

Procedure

1 [24 libraries] Arrange the index adapters in the TruSeq Index Plate Fixture as follows.

- Arrange Index 1 (i7) adapters in columns 1–6 of the TruSeq Index Plate Fixture.
- Arrange Index 2 (i5) adapter in rows A–D of the TruSeq Index Plate Fixture.

3 Using a multichannel pipette, add 5 µl of each Index 1 (i7) adapter down each column. Replace the cap on each i7 adapter tube with a new orange cap.

4 Using a multichannel pipette, add 5 µl of each Index 2 (i5) adapter across each row. Replace the cap on each i5 adapter tube with a new white cap.

5 Add 15 µl NPM to each well containing index adapters. Pipette to mix.

6 Centrifuge at 280 × g at 20°C for 1 minute.

7 Place on the preprogrammed thermal cycler and run the PCR program. The volume is 50 µl.

SAFE STOPPING POINT

If you are stopping, seal the plate and store at 2°C to 8°C for up to 2 days. Alternatively, leave on the thermal cycler overnight.

Clean Up Libraries

This step uses HighPrep™ PCR beads to purify the library DNA and remove short library fragments.

Consumables

- RSB (Resuspension Buffer)
- HighPrep™ PCR beads
- Freshly prepared 80% ethanol (EtOH)
- 96-well midi plate
- Hard-Shell 96-well PCR plate, skirted

About Reagents

- The HighPrep™ PCR beads are a user-supplied consumable.
- Vortex HighPrep™ PCR beads before each use.
- Vortex HighPrep™ PCR beads frequently to make sure that beads are evenly distributed.
- Always prepare fresh 80% ethanol for wash steps. Ethanol can absorb water from the air, impacting your results.

Preparation

1 Prepare the following consumables:

Item Storage Instructions
RSB -25°C to -15°C Thaw at room temperature.
RSB can be stored at 2°C to 8°C after the initial thaw.
HighPrep™ PCR Beads 2°C to 8°C Let stand on the benchtop for 30 minutes to bring to room temperature.

2 Prepare fresh 80% ethanol from absolute ethanol.

Procedure

1 Centrifuge at 280 × g at 20°C for 1 minute.

2 Transfer 50 µl PCR product from each well of the PCR plate to corresponding wells of a new midi plate.
NOTE: The ratio of PCR product to volume of beads is 3:2. For example, 50 µl PCR product to 30 µl AMPure. If you pull less than 50 µl of PCR product, adjust your ratio of AMPure beads accordingly.

3 Add 30 µl HighPrep™ PCR beads to each well. Smaller amplicons in Nextera XT library preps typically yield smaller insert size ranges. To maximize recovery of smaller fragments from the bead cleanup step, use the following conditions.

Input Size (bp) HighPrep™ PCR Recommendation HighPrep™ PCR Volume (µl)
300–500 1.8x HighPrep™ PCR 90
> 500 0.6x HighPrep™ PCR
(0.5x HighPrep™ PCR for ≥ 2 x 250
cycles)*
30
(25 µl for ≥ 2 x 250 cycles)*
gDNA or other genomic input 0.6x HighPrep™ PCR 30
*Applicable only to the MiSeq™ or HiSeq™ 2500 using HiSeq Rapid v2 reagents.

4 Shake at 1800 rpm for 2 minutes.
5 Incubate at room temperature for 5 minutes.
6 Place on a magnetic stand and wait until the liquid is clear (~2 minutes).
7 Remove and discard all supernatant from each well.
8 Wash two times as follows.
    a Add 200 µl fresh 80% EtOH to each well.
    b Incubate on the magnetic stand for 30 seconds.
    c Remove and discard all supernatant from each well.
9 Using a 20 µl pipette, remove residual 80% EtOH from each well.
10 Air-dry on the magnetic stand for 15 minutes.
11 Remove from the magnetic stand.
12 Add 52.5 µl RSB to each well.
13 Shake at 1800 rpm for 2 minutes.
14 Incubate at room temperature for 2 minutes.
15 Place on a magnetic stand and wait until the liquid is clear (~2 minutes).
16 Transfer 50 µl supernatant from the midi plate to a new Hard-Shell PCR plate.
SAFE STOPPING POINT
If you are stopping, seal the plate and store at -25°C to -15°C for up to seven days.

Check Libraries

1 Run 1 µl of undiluted library on an Agilent Technology 2100 Bioanalyzer using a High Sensitivity DNA chip.

Normalize Libraries

This process normalizes the quantity of each library to ensure more equal library representation in the pooled library.

Consumables

- LNA1 (Library Normalization Additives 1)
- LNB1 (Library Normalization Beads 1)
- LNW1 (Library Normalization Wash 1)
- LNS1 (Library Normalization Storage Buffer 1)
- 0.1 N NaOH (fewer than 7 days old) (3 ml per 96 samples)
- 96-well midi plate
- Hard-Shell 96-well PCR plate, skirted
- 15 ml conical tube
- Microseal 'B' adhesive seals

About Reagents

- Vortex LNA1 vigorously to make sure that all precipitates have dissolved. Inspect in front of a light.
- Vortex LNB1 vigorously, with intermittent inversion (at least 1 minute). Repeat until all beads are resuspended and no beads are present at the bottom of the tube when it is inverted.
- Always use a wide-bore pipette tip for LNA1.
- Mix only the required amounts of LNA1 and LNB1 for the current experiment. Store the remaining LNA1 and LNB1 separately at the recommended temperatures.
- Aspirate and dispense beads slowly due to the viscosity of the solution.

WARNING
This set of reagents contains potentially hazardous chemicals. Personal injury can occur through inhalation, ingestion, skin contact, and eye contact. Wear protective equipment, including eye protection, gloves, and laboratory coat appropriate for risk of exposure. Handle used reagents as chemical waste and discard in accordance with applicable regional, national, and local laws and regulations. For additional environmental, health, and safety information, see the SDS at support.illumina.com/sds.html.

Preparation

1 Prepare the following consumables:

Item Storage Instructions
LNA1 -25°C to -15°C Prepare under a fume hood.
Bring to room temperature. Use a 20°C to 25°C water bath as needed.
LNB1 2°C to 8°C Bring to room temperature. Use a 20°C to 25°C water bath as needed.
LNW1 2°C to 8°C Bring to room temperature. Use a 20°C to 25°C water bath as needed.
LNS1 Room temperature Bring to room temperature.

Procedure

1 Transfer 20 µl supernatant from the Hard-Shell PCR plate to a new midi plate.
2 Add 44 µl LNA1 per sample to a new 15 ml conical tube. Calculate about 5% extra sample to account for sample loss due to pipetting. For example: for 96 samples, add 4.4 ml LNA1 to the tube (100 samples × 44 µl = 4.4 ml).
3 Thoroughly resuspend LNB1. Pipette to mix.
4 Transfer 8 µl LNB1 per sample (including the 5% extra) to the 15 ml conical tube containing LNA1. Invert to mix. For example: for 96 samples, transfer 800 µl LNB1 to the tube of LNA1 (100 samples × 8 µl = 800 µl).
5 Pour the bead mixture into a trough.
6 Add 45 µl combined LNA1 and LNB1 to each well containing libraries.
7 Shake at 1800 rpm for 30 minutes.
8 Place on a magnetic stand and wait until the liquid is clear (~2 minutes).
9 Remove and discard all supernatant from each well.
10 Wash two times as follows.
    a Add 45 µl LNW1 to each well.
    b Shake at 1800 rpm for 5 minutes.
    c Place on a magnetic stand and wait until the liquid is clear (~2 minutes).
    d Remove and discard all supernatant from each well.
11 Add 30 µl 0.1 N NaOH to each well.
12 Shake at 1800 rpm for 5 minutes.
13 During the 5 minute elution, label a new 96-well PCR plate SGP for storage plate.
14 Add 30 µl LNS1 to each well of the SGP plate. Set aside.
15 After the 5 minute elution, make sure that all samples in the midi plate are resuspended. If they are not, resuspend as follows.
    a Pipette to mix or lightly tap the plate on the bench.
    b Shake at 1800 rpm for 5 minutes.
16 Place on a magnetic stand and wait until the liquid is clear (~2 minutes).
17 Transfer the supernatant from the midi plate to the SGP plate.
18 Centrifuge at 1000 × g for 1 minute.

NOTE
After denaturation, the libraries are single-stranded DNA, which resolves poorly on an agarose gel or Bioanalyzer chip. For quality control, use the double-stranded DNA saved from step 16 of the cleanup procedure.

SAFE STOPPING POINT
If you are stopping, seal the plate and store at -25°C to -15°C for up to seven days.

Pool Libraries

Pooling libraries combines equal volumes of normalized libraries in a single tube. After pooling, dilute and heat-denature the library pool before loading libraries for the sequencing run.

Consumables

- Adhesive PCR foil seal
- Eppendorf LoBind microcentrifuge tubes
- PCR eight-tube strip

Preparation

1 To prepare for the sequencing run, begin thawing reagents according to the instructions for your instrument.
2 If the SGP plate was stored frozen at -25°C to -15°C, thaw at room temperature. Pipette to mix.

Procedure

1 Centrifuge at 1000 × g at 20°C for 1 minute.
2 Label a new Eppendorf tube PAL.
3 Transfer 5 µl of each library from the SGP plate to the PAL tube. Invert to mix.
4 Dilute pooled libraries to the loading concentration for your sequencing system. For instructions, see the denature and dilute libraries guide for your system.
5 Store unused pooled libraries in the PAL tube and SGP plate at -25°C to -15°C for up to 7 days.

S7: Overview of costs for "in-house" library prep based either on a commercial kit (Nextera XT) or on two custom-made pipelines. Results are also reported for outsourcing to Macrogen (96 samples) and MGI Sequencing Technology (227 samples).

JEMU Nextera XT DNA Library Preparation Kit		
n. samples		96
Nextera XT DNA Library Preparation Kit (+VAT)	€	4,431
TG Nextera® XT Index Kit v2 Set B (96 Indices) (+VAT)	€	1,479
library QC Genomics Core KUL	€	50
Sequencing - MiSeq - 300 PE (1 FC)	€	1,900
cost / plate	€	7,956
cost / sample (personnel costs @60k €/yr, gross, 1 working week)	€	1,154
cost / sample (consumables)	€	83
cost / sample (consumables + personnel)	€	95

JEMU semi-custom library prep		
n. samples		96
TG Nextera® XT Index Kit v2 Set B (96 Indices) (+VAT)	€	1,479
custom primers with hoverhangs adapters	€	110
Agencourt AMPure XP (full cost)	€	516
library QC Genomics Core KUL	€	50
Sequencing - MiSeq - 300 PE (1 FC)	€	1,900
	€	4,151
	€	1,154
cost / sample (consumables)	€	43
cost / sample (consumables + personnel)	€	55

JEMU fully-custom library prep		
n. samples		96
custom primers with hoverhangs adapters	€	110
Agencourt AMPure XP (full cost)	€	516
Custom indices with Illumina adapters (full cost)	€	462
library QC Genomics Core KUL	€	50
Sequencing - MiSeq - 300 PE (1 FC)	€	1,900
	€	3,134
	€	1,154
cost / sample (consumables)	€	33
cost / sample (consumables + personnel)	€	45

Outsourced Macrogen		
n. samples		96
metabarcoding library prep	€	2,112
Sequencing - MiSeq - 300 PE (1 FC)	€	1,900
	€	4,108
cost / sample (consumables + personnel)	€	43

Outsourced MGI Sequencing Technology (best offer RMCA)		
n. samples		227
metabarcoding library prep + Sequencing - MiSeq - 300 PE (1 FC)	€	5,000
cost / sample (consumables + personnel)	€	22